arXiv:1210.5127v1 [math.CV] 18 Oct 2012

AN ENTIRE FUNCTION WITH NO FIXED POINTS AND NO INVARIANT BAKER DOMAINS

WALTER BERGWEILER

Abstract. We show that there exists an entire function which has neither fixed points nor invariant Baker domains. The question whether such a function exists was raised by Buff.

1. Introduction and result

Let f be a meromorphic function and denote by f^n the n-th iterate of f. An invariant component U of the Fatou set of f such that $f^n|_U \to \infty$ as $n \to \infty$ is called an *invariant Baker domain*; cf. [3, §4.7] or [11]. It was suggested by Douady that invariant Baker domains of the Newton function $f(z) = z - g(z)/g'(z)$ of an entire function g are related to paths where g tends to the asymptotic value 0. In response to Douady's question it was shown in [8] that under mild additional hypotheses the existence of an invariant Baker domain does indeed imply that 0 is an asymptotic value of g. However, this is not always the case [5].

If g has no zeros at all, then the Newton function f has no fixed points. Moreover, 0 is an asymptotic value of g by Iversen's theorem [10, p. 289]. This led Buff to ask whether there exists an entire function having no fixed points and no invariant Baker domains. We show that such a function exists.

Theorem. *There exists an entire function with no fixed points and no invariant Baker domains.*

A meromorphic function with this property was constructed in [4]. The present construction is based on similar ideas. As in [4], a function f satisfying the conclusion of the theorem can be given explicitly.

Let (r_k) be a sequence of real numbers tending to ∞ and let (n_k) be a sequence of positive integers satisfying $n_k \geq k$ for all $k \in \mathbb{N}$. Then

$$h(z) = \prod_{k=1}^{\infty} \left(1 + \left(\frac{z}{r_k}\right)^{n_k}\right)$$

defines an entire function h. Indeed, if $|z| \leq R$ and k is so large that $r_k \geq 2R$, then $|z/r_k|^{n_k} \leq 2^{-k}$, implying that the infinite product converges locally uniformly.

For $k \geq 2$ we put $m_k = \sum_{j=1}^{k-1} n_j$. We shall show that if

$$r_k \geq 2r_{k-1} \geq 4 \quad \text{and} \quad n_k \geq 20 r_k^2 \exp\left(4 r_k^{m_k}\right) \tag{1.1}$$

for $k \geq 2$, then $f(z) = z + e^{h(z)}$ has the required property.

Acknowledgment. I thank Alexandre Eremenko for useful comments.

2010 *Mathematics Subject Classification.* Primary 37F10; Secondary 30D05, 49M15, 65H05.

Supported by the ESF Research Networking Programme HCAA.

2. Preliminaries

2.1. **The hyberbolic metric.** We need some standard results about the hyperbolic metric which can be found in, e.g., [9, Section I.4].

We denote the open disk of radius r around a point $c \in \mathbb{C}$ by $D(c,r)$ and put $\mathbb{D} = D(0,1)$. The density of the hyperbolic metric in a hyperbolic domain U is denoted by λ_U, normalized such that $\lambda_{\mathbb{D}}(z) = 2/(1-|z|^2)$. The hyperbolic metric is denoted by ρ_U. For $a, b \in U$ we thus have

$$\rho_U(a,b) = \inf_\gamma \int_\gamma \lambda_U(z)|dz|,$$

where the infimum is taken over all curves γ that connect a and b. Then [9, p. 11]

$$\rho_{\mathbb{D}}(0,z) = \log \frac{1+|z|}{1-|z|} \quad \text{for } z \in \mathbb{D}. \tag{2.1}$$

It follows from Schwarz's lemma and the Koebe one quarter theorem that if U is simply connected, then [9, Theorem I.4.3]

$$\frac{1}{2\operatorname{dist}(z,\partial U)} \le \lambda_U(z) \le \frac{2}{\operatorname{dist}(z,\partial U)} \tag{2.2}$$

for all $z \in U$. Here $\operatorname{dist}(z,\partial U) = \inf_{\zeta\in\partial U} |\zeta - z|$.

The following lemma is a simple consequence of (2.2).

Lemma 1. *Let U be a simply connected hyperbolic domain, $a, b \in U$ and $c \in \mathbb{C}\backslash U$. Then*

$$\rho_U(a,b) \ge \frac{1}{2}\left|\log\left|\frac{b-c}{a-c}\right|\right|.$$

Proof. Without loss of generality we may assume that $c = 0$. Let γ be a curve from a to b and let L be a branch of the logarithm defined in U. Then (2.2) yields

$$\begin{aligned}\int_\gamma \lambda_U(z)|dz| &\ge \frac{1}{2}\int_\gamma \frac{|dz|}{\operatorname{dist}(z,\partial U)} \ge \frac{1}{2}\int_\gamma \frac{|dz|}{|z|} \ge \frac{1}{2}\left|\int_\gamma \frac{dz}{z}\right| \\ &= \frac{1}{2}|L(b)-L(a)| \ge \frac{1}{2}|\operatorname{Re}(L(b)-L(a))| = \frac{1}{2}\left|\log\left|\frac{b}{a}\right|\right|,\end{aligned}$$

from which the conclusion follows. ☐

The next lemma follows easily from (2.1) and the triangle inequality.

Lemma 2. *If $a, b \in D(c, r/2)$, then $\rho_{D(c,r)}(a,b) \le 2\log 3$.*

Finally we have the following form of Schwarz's lemma [9, Theorem I.4.3].

Lemma 3. *Let U, V be hyperbolic domains, $f \colon U \to V$ holomorphic and $a, b \in U$. Then $\rho_V(f(a), f(b)) \le \rho_U(a,b)$.*

Applying this lemma to $f(z) = z$ yields

$$\rho_V(a,b) \le \rho_U(a,b) \quad \text{if } U \subset V. \tag{2.3}$$

2.2. **Some growth estimates.** We have to estimate the growth of h on certain circles from above and below. For $k \geq 2$ and $|z| = r_k$ we have

$$\begin{aligned}
\log|h(z)| &\leq \sum_{j=1}^{k-1} \log\left(1+\left(\frac{r_k}{r_j}\right)^{n_j}\right) + \log 2 + \sum_{j=k+1}^{\infty} \log\left(1+\left(\frac{r_k}{r_j}\right)^{n_j}\right) \\
&\leq \sum_{j=1}^{k-1} \log\left(1+\frac{1}{2}{r_k}^{n_j}\right) + \log 2 + \sum_{j=k+1}^{\infty} \left(\frac{r_k}{r_j}\right)^{n_j} \\
&\leq \sum_{j=1}^{k-1} \log\left({r_k}^{n_j}\right) + \log 2 + \sum_{j=k+1}^{\infty} 2^{-n_j} \leq m_k \log r_k + 2\log 2.
\end{aligned}$$

Hence

$$|h(z)| \leq 4 r_k^{m_k} \quad \text{for } |z| = r_k \tag{2.4}$$

and $k \geq 2$.

We put $s_k = (1 + 1/n_k) r_k$. For $t \in [0, 2\pi]$ and $z = s_k e^{it}$ we have

$$\begin{aligned}
h(z) &= \prod_{j=1}^{k-1}\left(1+\left(\frac{z}{r_j}\right)^{n_j}\right) \cdot \left(1+\left(1+\frac{1}{n_k}\right)^{n_k} e^{in_k t}\right) \cdot \prod_{j=k+1}^{\infty}\left(1+\left(\frac{z}{r_j}\right)^{n_j}\right) \\
&\sim \prod_{j=1}^{k-1}\left(\frac{z}{r_j}\right)^{n_j}\left(1+e\cdot e^{in_k t}\right) = \prod_{j=1}^{k-1}\left(\frac{s_k}{r_j}\right)^{n_j} e^{im_k t}\left(1+e\cdot e^{in_k t}\right)
\end{aligned}$$

as $k \to \infty$. Putting

$$T_k = \prod_{j=1}^{k-1}\left(\frac{s_k}{r_j}\right)^{n_j} \tag{2.5}$$

we thus have

$$h(s_k e^{it}) \sim T_k e^{im_k t}\left(1+e\cdot e^{in_k t}\right) \tag{2.6}$$

as $k \to \infty$.

It is not difficult to see that for each $\varphi \in \mathbb{R}$ there exists $\theta = \theta(\varphi) \in [0,1]$ such that $e^{2\pi i\varphi}\left(1+e\cdot e^{2\pi i\theta}\right)$ is positive. For $\nu \in \{0, 1, \ldots, n_k - 1\}$ we put

$$\theta_\nu = \theta(\nu m_k/n_k) \quad \text{and} \quad p_\nu = e^{2\pi i \nu m_k/n_k}\left(1+e\cdot e^{2\pi i\theta_\nu}\right).$$

Then p_ν is positive and thus $p_\nu = |p_\nu| \geq e - 1$. Since $m_k/n_k \to 0$ by (1.1), we deduce from (2.6) that

$$h(s_k e^{2\pi i(\nu+\theta_\nu)/n_k}) \sim T_k e^{2\pi i(\nu+\theta_\nu)m_k/n_k}\left(1+e\cdot e^{2\pi i(\nu+\theta_\nu)}\right) = p_\nu T_k e^{2\pi i\theta_\nu m_k/n_k} \sim p_\nu T_k$$

Thus

$$\operatorname{Re} h(s_k e^{2\pi i(\nu+\theta_\nu)/n_k}) \geq T_k \tag{2.7}$$

for large k and all $\nu \in \{0, 1, \ldots, n_k - 1\}$.

3. PROOF OF THE THEOREM

Let f be as defined in the introduction and suppose that f has an invariant Baker domain U. By a result of Baker [1], U is simply connected. Take $z_0 \in U$, connect z_0 and $f(z_0)$ by a curve γ_0 in U and put $\gamma = \bigcup_{j=0}^{\infty} f^j(\gamma_0)$. Then γ is a curve in U connecting z_0 to ∞. As γ_0 is compact, there exists $K > 0$ such that $\rho(f(z), z) \leq K$ for all $z \in \gamma_0$. Since every $z \in \gamma$ has the form $z = f^j(\zeta)$ for some $\zeta \in \gamma_0$ and some $j \geq 0$, Lemma 3 yields

$$\rho(f(z), z) \leq K \quad \text{for } z \in \gamma. \tag{3.1}$$

For large k the curve γ intersects the circle $\{z \colon |z| = r_k\}$. Let z_k be a point of intersection.

We shall show first that if k is large enough, then the disk $D(z_k, 20r_k/n_k)$ is not contained in U; that is,

$$D(z_k, 20r_k/n_k) \cap \partial U \neq \emptyset. \tag{3.2}$$

In order to do so we assume that $D(z_k, 20r_k/n_k) \subset U$. We write $z_k = r_k e^{2\pi i t_k}$ with $t_k \in [0, 1)$ and put $\nu = [n_k t_k]$, where $[x]$ denotes the largest integer not greater than x. Thus $n_k t_k = \nu + \delta$ where $\nu \in \{0, 1, \ldots, n_k - 1\}$ and $\delta \in [0, 1)$. Let

$$a_k = r_k e^{(2\nu+1)\pi i/n_k} \quad \text{and} \quad b_k = s_k e^{2\pi i(\nu+\theta_\nu)/n_k}.$$

Then

$$|a_k - z_k| = r_k \left| e^{(2\nu+1)\pi i/n_k - 2\pi i t_k} - 1 \right| = r_k \left| e^{(1-2\delta)\pi i/n_k} - 1 \right| \sim \frac{|1-2\delta|\pi r_k}{n_k}$$

and

$$\begin{aligned} |b_k - z_k| &\leq |b_k - r_k e^{2\pi i(\nu+\theta_\nu)/n_k}| + |r_k e^{2\pi i(\nu+\theta_\nu)/n_k} - z_k| \\ &= s_k - r_k + r_k \left| e^{2\pi i(\nu+\theta_\nu)/n_k - 2\pi i t_k} - 1 \right| \\ &= \frac{r_k}{n_k} + r_k \left| e^{2\pi i(\theta_\nu - \delta)/n_k} - 1 \right| \\ &\sim \frac{(1 + 2\pi|\theta_\nu - \delta|) r_k}{n_k}, \end{aligned}$$

which implies that

$$a_k \in D(z_k, 10r_k/n_k) \quad \text{and} \quad b_k \in D(z_k, 10r_k/n_k)$$

for large k. Lemma 2 and (2.3) now yield

$$\rho_U(a_k, b_k) \leq \rho_{D(z_k, 20r_k/n_k)}(a_k, b_k) \leq 2\log 3. \tag{3.3}$$

Since $h(a_k) = 0$ by the definition of h and $\operatorname{Re} h(b_k) \geq T_k \geq s_k/r_1$ by (2.5) and (2.7), we have

$$|f(a_k)| = |a_k + 1| \leq r_k + 1 \quad \text{and} \quad |f(b_k)| \geq e^{s_k/r_1} - s_k \geq s_k^2 \geq r_k^2 \tag{3.4}$$

for large k. Fix a point $c \in \partial U$. Lemma 1 and (3.4) imply that

$$\rho_U(f(a_k), f(b_k)) \geq \frac{1}{2} \log \left| \frac{f(b_k) - c}{f(a_k) - c} \right| \geq \frac{1}{2} \log \frac{r_k^2 - |c|}{r_k + 1 + |c|} \tag{3.5}$$

for large k. Now a contradiction is obtained from Lemma 3, (3.3) and (3.5), provided k is sufficiently large. This contradiction shows that (3.2) holds for large k.

Thus, for large k, there exists $c_k \in D(z_k, 20r_k/n_k) \cap \partial U$. Lemma 1 now yields

$$\rho_U(f(z_k), z_k) \geq \frac{1}{2} \log \left| \frac{f(z_k) - c_k}{z_k - c_k} \right| = \frac{1}{2} \log \left| \frac{e^{h(z_k)}}{z_k - c_k} + 1 \right| .$$

Since

$$\left| \frac{e^{h(z_k)}}{z_k - c_k} \right| \geq \frac{e^{-|h(z_k)|}}{|z_k - c_k|} \geq \frac{n_k \exp\left(-4r_k^{m_k}\right)}{20r_k} \geq r_k$$

for $k \geq 2$ by (1.1) and (2.4), we obtain

$$\rho_U(f(z_k), z_k) \geq \frac{1}{2} \log(r_k - 1)$$

for large k, contradicting (3.1). ☐

Remark 1. Buff and Rückert [8] considered *virtual immediate basins* instead of invariant Baker domains. However, for functions for which all Baker domains are simply connected the two concepts coincide; cf. the discussion in [4, p. 431] or [8, p. 4]. By the result of Baker [1] already quoted, this holds in particular for entire functions. By a recent result of Barański, Fagella, Jarque and Karpińska [2], it also holds for Newton maps of entire functions.

Remark 2. The function f constructed in the proof is the Newton function for

$$g(z) = \exp\left(- \int_0^z e^{-h(t)} dt \right) .$$

Since g has no zeros, g has a direct singularity over 0; see [10, p. 289] for this result, as well as [6, 12] for the terminology used here and below. As f has no invariant Baker domains, g has no logarithmic singularity over 0 by one of the results obtained by Buff and Rückert in the paper already mentioned in the introduction [8, Theorem 4.1]. Thus g has a direct non-logarithmic singularity over 0. This implies [7, Theorem 5] that g has uncountably many direct non-logarithmic singularities over 0. As g has no critical points, a result of Sixsmith [12, Theorem 1.2] yields that every neighborhood of any of these singularities contains a neighborhood of an indirect or logarithmic singularity of g whose projection is different from 0. Overall we see that the set of singularities of the inverse of g has a quite complicated structure.

REFERENCES

[1] I. N. Baker, The domains of normality of an entire function. Ann. Acad. Sci. Fenn. Ser. A I Math. 1 (1975), 277–283.

[2] Krzysztof Barański, Núria Fagella, Xavier Jarque and Bogusława Karpińska, On the connectivity of the Julia sets of meromorphic functions. Preprint, arXiv: 1206.6667.

[3] Walter Bergweiler, Iteration of meromorphic functions. Bull. Amer. Math. Soc. (N. S.) 29 (1993), 151–188.

[4] Walter Bergweiler, Newton's method and Baker domains. J. Difference Equ. Appl. 16 (2010), 427–432.

[5] Walter Bergweiler, David Drasin and James K. Langley, Baker domains for Newton's method. Ann. Inst. Fourier 57 (2007), 803–814.

[6] Walter Bergweiler and Alexandre Eremenko, On the singularities of the inverse to a meromorphic function of finite order. Rev. Mat. Iberoamericana 11 (1995), 355–373.

[7] Walter Bergweiler and Alexandre Eremenko, Direct singularities and completely invariant domains of entire functions. Illinois J. Math. 52 (2008), 243–259.

[8] Xavier Buff and Johannes Rückert, Virtual immediate basins of Newton maps and asymptotic values. Int. Math. Res. Not. 2006, 65498, 1–18.

[9] Lennart Carleson and Theodore W. Gamelin, Complex Dynamics. Universitext: Tracts in Mathematics. Springer-Verlag, New York, 1993.

[10] Rolf Nevanlinna, Eindeutige analytische Funktionen. Springer-Verlag, Berlin, Göttingen, Heidelberg, 1953.

[11] P. J. Rippon, Baker domains. In "Transcendental Dynamics and Complex Analysis". London Math. Soc. Lect. Note Ser. 348. Edited by P. J. Rippon and G. M. Stallard, Cambridge University Press, Cambridge, 2008, 371–395.

[12] D. J. Sixsmith, A new characterisation of the Eremenko-Lyubich class. Preprint, arXiv: 1205.1976.

Mathematisches Seminar, Christian–Albrechts–Universität zu Kiel, Ludewig–Meyn–Str. 4, D–24098 Kiel, Germany

E-mail address: `bergweiler@math.uni-kiel.de`